US ERA ARCHIVE DOCUMENT

Air Toxics in the Detroit Exposure and Aerosol Research Study

Alan Vette, Donald Whitaker, Timothy Barzyk, BJ George, Bob Seila, Shaibal Mukerjee, and Ron Williams


U.S. EPA

National Exposure Research Laboratory

Detroit, MI October 24, 2007

DEARS Study Sites

(Emphasis on Proximity to Sources)







DEARS Monitoring Design

- 3 year study July '04 to Feb. '07
- ~120 homes 5 days in winter and summer
- Concurrent monitoring
 - Personal
 - Residential indoor & outdoor
 - Ambient






DEARS Measurements

<u>Parameter</u>	Personal	<u>Indoor</u>	Outdoor	<u>Ambient</u>
PM _{2.5} (mass, elements)	X	x	х	x
PM _{coarse} (mass, elements)		X	х	х
EC-OC (PM _{2.5})		X	X	X
EC (PM _{2.5})	X	x	Х	х
Nitrate		x	х	x
Gases (O ₃ , NO ₂ , SO ₂)	Х		-NO ₂	x
Aldehydes	Х	х	х	x
VOCs	Х	х	X	x
SVOCs		x	Х	x
PAHs		x	X	х
Air Exchange Rate		X		



Passive Samplers





- Diffusive samplers -25 VOCs and 3 aldehydes
- Aromatics/HCs (9)
- Halogenated HCs (16)
- Detection limits
 - VOCs ~ 50 pptv



VOCs Measured in DEARS

Aromatics/HCs

Benzene

Toluene

Ethylbenzene

m,p-Xylene

o-Xylene

1,3-Butadiene

4-Ethyltoluene

1,3,5-Trimethylbenzene

Styrene

Chlorinated Aromatics/HCs

Carbon tetrachloride

1,1-Dichloroethane

1,2-Dichloroethane

1,1,1-Trichloroethane

1,1-Dichloroethylene

cis-1,2-Dichloroethylene

Trichloroethylene

Perchloroethylene

1,2-Dichloropropane

Chlorobenzene

m-Dichlorobenzene

o-Dichlorobenzene

p-Dichlorobenzene


<u>Freon</u>

Trichlorofluoromethane (11)

1,1,2-Trichloro-1,2,2-trifluoroethane (113)


1,2-Dichloro-1,1,2,2tetrafluoroethane (114)






Preliminary results – Winter '06





Preliminary results – Winter '06





Preliminary results – Winter '06



Compound	Slope ^a	Intercept	r²
Benzene	1.01 ± 0.13	-0.61 ± 0.20 ^a	0.69
Toluene	0.92 ± 0.11	-0.65 ± 0.33^{b}	0.73
Ethylbenzene	0.62 ± 0.11	0.31 ± 0.05^{a}	0.57
o-Xylene	0.86 ± 0.11	0.17 ± 0.05^{a}	0.72
m,p-Xylenes	0.84 ± 0.13	0.60 ± 0.17 ^a	0.61
1,3-Butadiene	1.66 ± 0.27	-0.01 ± 0.04°	0.63

a p<0.01


 $^{\mathrm{b}}$ p<0.1

Preliminary results – Winter '06



^c Not significant (p >> 0.10)


Duplicate Tube SamplesOutdoor and Ambient



Preliminary results – Summer '04 – Winter '06




Duplicate Tube SamplesOutdoor and Ambient





Duplicate Tube SamplesOutdoor and Ambient



Preliminary results – Summer '04 – Winter '06



Duplicate SamplesOutdoor and Ambient

Compound	Slope ^a	Interceptb	r²
Benzene	0.93 ± 0.06	0.25 ± 0.19	0.76
Toluene	0.93 ± 0.05	0.45 ± 0.31	0.84
Ethylbenzene	1.10 ± 0.09	-0.01 ± 0.11	0.68
o-Xylene	1.13 ± 0.09	-0.01 ± 0.12	0.68
m,p-Xylenes	1.15 ± 0.11	-0.04 ± 0.43	0.68
1,3-Butadiene	0.94 ± 0.06	0.07 ± 0.05	0.76


a p<0.001

Preliminary results – Summer '04 – Winter '06




^b Not significant (p >> 0.10)

Impact of ETS




- Indoor levels of VOCs associated with tobacco smoke
- ETS was measured using an optical technique on filter samples
- Samples were considered impacted by ETS when ETS levels exceeded
 1.5 µg m⁻³








Concentration Ratios Outdoor Residential to Ambient Site






Preliminary results – Summer '04 – Summer '05

www.epa.gov/DEARS





Industrial

Personal Exposure Relationships



Benzene










Personal Exposure Relationships

1,3-Butadiene










Personal Exposure Relationships Log Transformed Data



- Mixed models account for repeated measures
- Exposures related to indoor, outdoor and sometimes ambient
- Slopes (agreement):I > O > A



Conclusions

- Carbopack-X passive tubes robust method to measure VOCs
- Second hand smoke found in 16% of indoor air and 22% of personal exposure samples
- Significant spatial and temporal variability in MSATs
 - Impact of sources (EMA) and locations (I, O, A)
- Personal exposures to MSATs
 - Indoor > Outdoor > Ambient
 - Concentrations and relationships



Disclaimer

Although this work was reviewed by EPA and approved for publication, it may not necessarily reflect official Agency policy.

The U.S. Environmental Protection Agency through its Office of Research and Development funded and conducted the research described here through contract 68-D-00-012 with RTI International, EP-D-04-068 to Battelle Columbus Laboratory, 68-D-00-206 and EP-05-D-065 to Alion Science and Technology.



